

MEMORANDUM REPORT ARBRL-MR-03335

CALCULATION OF LEGENDRE FUNCTIONS ON THE  
CUT FOR INTEGRAL ORDER AND COMPLEX  
DEGREE BY MEANS OF GAUSS  
CONTINUED FRACTIONS

Alexander S. Elder  
James N. Walbert  
Eric C. Benck

February 1984



US ARMY ARMAMENT RESEARCH AND DEVELOPMENT CENTER  
**BALLISTIC RESEARCH LABORATORY**  
ABERDEEN PROVING GROUND, MARYLAND

Approved for public release; distribution unlimited.

Destroy this report when it is no longer needed.  
Do not return it to the originator.

Additional copies of this report may be obtained  
from the National Technical Information Service,  
U. S. Department of Commerce, Springfield, Virginia  
22161.

The findings in this report are not to be construed as  
an official Department of the Army position, unless  
so designated by other authorized documents.

*The use of trade names or manufacturers' names in this report  
does not constitute indorsement of any commercial product.*

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Memorandum Report ARBRL-MR-03335	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  CALCULATION OF LEGENDRE FUNCTIONS ON THE CUT FOR INTEGRAL ORDER AND COMPLEX DEGREE BY MEANS OF GAUSS CONTINUED FRACTIONS		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) ALEXANDER S. ELDER JAMES N. WALBERT ERIC C. BENCK		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS US Army Ballistic Research Laboratory, ARDC ATTN: DRSMC-BLI(A) Aberdeen Proving Ground, MD 21005		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS US Army AMCCOM, ARDC Ballistic Research Laboratory, ATTN: DRSMC-BLA-S(A) Aberdeen Proving Ground, MD 21005		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS  1L162618AH80
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE February 1984
		13. NUMBER OF PAGES 49
		15. SECURITY CLASS. (of this report)  UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Legendre Functions ✓ Gauss Continued Fraction Continued Fraction <i>Math Models</i> <i>Bi-harmonic functions</i>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) jmk A method for calculating the associated Legendre functions for integral order and complex degree is presented. Comparison with special cases indicates precision of at least 25 significant digits is attained.		

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

## TABLE OF CONTENTS

	PAGE
I. INTRODUCTION.....	5
II. ASSOCIATED LEGENDRE FUNCTIONS OF THE FIRST KIND.....	6
A. Hypergeometric Series.....	6
B. Gauss Contiguous Formula.....	7
III. ASSOCIATED LEGENDRE FUNCTIONS OF THE SECOND KIND.....	8
A. Gauss Continued Fraction Method.....	8
B. The Logarithmic Solution.....	13
IV. CONCLUSION.....	15
REFERENCES.....	17
APPENDIX A. NUMERICAL CALCULATIONS.....	19
APPENDIX B. GRAPHS. ....	35
DISTRIBUTION LIST.....	45

## I. INTRODUCTION

The associated Legendre functions on the cut for integral order and complex degree are required for several important mechanical applications. One area of particular interest is in the calculation of biharmonic functions for the hollow cone. Such functions appropriate for the cone satisfy

$$\nabla^4 \chi = 0$$

and come in the form

$$\chi = [AP_v^m(\cos\theta) + BQ_v^m(\cos\theta) + CP_{v-2}^m(\cos\theta) + DQ_{v-2}^m(\cos\theta)] R^v \sin m\phi.$$

These functions are required for stress analysis in the mini-hat gage, which is a strain-type pressure transducer, and in cone-plate rheometers.

We have developed a subroutine for computing

$$P_v^m(\cos\theta) \text{ and } Q_v^m(\cos\theta)$$

for

$$0 < \theta < 90^\circ,$$

$$\text{integer } m \text{ such that } 0 \leq m \leq 12,$$

$$\text{and } v \text{ such that } \operatorname{Re}(v) > -1/2.$$

For moderate values of  $|v|$ , we calculate  $P_v^m(\cos\theta)$  by series and recurrence formulas. On the other hand,  $Q_v^m(\cos\theta)$  is computed through the use of a Gauss continued fraction and Wronskian relations. For small angles, the logarithmic solution is used to calculate  $Q_v^m(\cos\theta)$ . Additional analysis will be

necessary for the programming of  $\frac{\partial Q_v^m(\cos\theta)}{\partial v}$  and  $\frac{\partial P_v^m(\cos\theta)}{\partial v}$ . Asymptotic expansions are required when  $|v|$  is large.

Programming was done on a CDC 7600 computer in FORTRAN V. CDC double precision arithmetic was used to get the greatest accuracy possible. Special

multiple precision algorithms were avoided so that the subroutine would be more compatible for use by an engineering laboratory.

## II. ASSOCIATED LEGENDRE FUNCTIONS OF THE FIRST KIND

### A. Hypergeometric Series

The Legendre functions of the first kind can be accurately calculated for moderate  $|v|$  by<sup>1</sup>

$$P_v^{-m}(\cos\theta) = \frac{1}{m!} (\tan \theta/2)^m {}_2F_1(-v, v+1; l+m; (\sin\theta/2)^2) \quad (1)$$

and the reflection formula<sup>2</sup>

$$P_v^m(\cos\theta) = (-1)^m \frac{\Gamma(v+m+1)}{\Gamma(v-m+1)} P_v^{-m}(\cos\theta), \quad (2)$$

where

$$\frac{\Gamma(v+m+1)}{\Gamma(v-m+1)} = (v+1-m)(v+2-m)\dots(v+m). \quad (2a)$$

The hypergeometric series, for small  $|v|$ , ( $|v| < 2$ ), can be calculated directly from the series<sup>3</sup>

$${}_2F_1(a, b; c; z) = 1 + \frac{ab}{c} \frac{z}{1!} + \frac{a(a+1)b(b+1)}{c(c+1)} \frac{z^2}{2!} + \dots \quad (3)$$

For larger values of  $|v|$ , the series becomes slowly convergent with serious round-off error. In this case, one can alter the parameters and then use a recurrence formula to obtain the desired results.

<sup>1</sup> A. Erdelyi, et. al., *Higher Transcendental Functions*, Vol. 1, Bateman Manuscript Project, McGraw Hill Book Co., Inc., New York, 1953.

<sup>2</sup> W. Magnus and F. Oberhettinger, *Formulas and Theorems for the Special Functions of Mathematical Physics*, Chelsea Publishing Co., New York, 1949.

<sup>3</sup> M. Abramowitz and J.A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Washington, D.C., No. 55, Applied Mathematics Series, June 1964.

## B. Gauss Contiguous Formula

One possible method to improve the convergence of the hypergeometric series would be to increase  $m$  to a high value and then use a Legendre recurrence formula to obtain the desired answer. Unfortunately, either serious underflow or overflow problems would begin to occur in the calculation of the initial values for the Legendre recurrence formula, as found by Smith, Olver, and Lozier.<sup>4</sup> Occasionally both underflow and overflow occur in the same sequence of calculations. The necessary recurrence formula,<sup>5</sup>

$$P_v^{m+1}(\cos\theta) + 2m\cot\theta P_v^m(\cos\theta) + (v+m)(v-m+1)P_v^{m-1}(\cos\theta) = 0, \quad (4)$$

also contains both linear and quadratic coefficients which would lead to rapid variations of the Legendre functions in a recurrence chain.

In order to avoid the difficulties in dealing with the Legendre function, we deal directly with the hypergeometric series using the Gauss contiguous formula. In the required Gauss contiguous formula,<sup>1</sup>

$$c(c-1)(z-1)F(a,b;c-1;z) + c[c-1-(2c-a-b-1)z]F(a,b;c;z) + (c-a)(c-b)z F(a,b;c+1;z) = 0, \quad (5)$$

the coefficients of the hypergeometric functions are of the same degree in the parameter  $c$ . Therefore, the values of the hypergeometric functions vary slowly as the parameters change in the recurrence chain. Also, the stability criterion for the Gauss contiguous formula can be verified by elementary methods. Finally, through the use of the hypergeometric functions as the basis of our analysis, we gain an additional parameter leading to greater flexibility in deriving algorithms.

Since the Gauss contiguous formula appears to be superior for the intended work, it is used in the program. We increase  $c$  to a sufficiently large value and then increment downwards with the Gauss contiguous formula to the desired result. Using this technique has produced excellent results, provided that  $|v|$  is not too large.

---

<sup>4</sup> J.M. Smith, F.V.J. Olver and D.W. Lozier, "Legendre Polynomials," *ACM Transactions on Mathematical Software*, 7, 94-95, 1981.

<sup>5</sup> A.S. Elder, "Formulas for Calculating Bessel Functions of Integral Order and Complex Argument," *USA Ballistic Research Laboratory, Report No. BRL-TR-1423*, 1968 AD #680209.



### III. ASSOCIATED LEGENDRE FUNCTIONS OF THE SECOND KIND

#### A. Gauss Continued Fraction Method

In attempting to calculate  $Q_v^m(\cos\theta)$ , no series was found which converges well for small  $v$  and  $\theta$ . Instead, an approach previously used to calculate Bessel functions of the second kind was used.<sup>5</sup> This approach utilizes the Gauss continued fraction and Wronskian relations. It was found to be highly advantageous to leave the cut for the initial calculations and work in the complex plane. However,  $Q_v^m(x)$ , as defined on the cut  $-1 < x < +1$ , is not analytic when  $x=z$ , a complex number. We have the formula<sup>1</sup>

$$Q_v^m(x) = \frac{1}{2} e^{-im\pi} \left[ e^{-\frac{1}{2} i\pi} Q_v^m(x+i0) + e^{\frac{1}{2} i\pi} Q_v^m(x-i0) \right], \quad (6)$$

where  $x+i0$ ,  $x-i0$  indicate limits as the complex variable  $z$  approaches the cut. If  $v = \alpha + i\beta$ , and  $\beta$  is large,

$$Q_v^m(\cos\theta+i0) \text{ is } O(e^{-\beta\theta})$$

and

$$Q_v^m(\cos\theta-i0) \text{ is } O(e^{+\beta\theta}), \text{ where } x = \cos\theta.$$

This behavior corresponds to that of Hankel functions in the theory of Bessel functions.

In the complex plane, we have<sup>1</sup>

$$\begin{aligned} e^{-im\pi} Q_v^m(\cos\theta+i0) \Gamma(v+3/2) &= \pi^{1/2} \Gamma(v+m+1) e^{\frac{1}{2} m\pi} (2\sin\theta)^m \\ &\times e^{-i\theta(1+v+m)} F\left(\frac{1}{2} + m, 1+v+m; v+3/2; e^{-2i\theta}\right). \end{aligned} \quad (7)$$

Replacing  $v$  by  $v-1$  and then dividing the result into the previous equation,

$$R_v^m(\cos\theta+i0) = \frac{Q_v^m(\cos\theta+i0)}{Q_{v-1}^m(\cos\theta+i0)} = \frac{(v+m)}{(v+\frac{1}{2})} e^{-i\theta} \frac{F\left(\frac{1}{2} + m, 1+v+m; v+3/2; e^{-2i\theta}\right)}{F\left(\frac{1}{2} + m, v+m; v+\frac{1}{2}; e^{-2i\theta}\right)}. \quad (8)$$

Letting  $a = 1/2 + m, b = v + m, c = v + 1/2, z = e^{-2i\theta}$ ,

$$R_v^m(\cos\theta + i0) = \frac{b e^{-i\theta}}{c} \frac{F(a, b+1; c+1; z)}{F(a, b; c; z)}. \quad (9)$$

This can be calculated by using the Gauss continued fraction<sup>6</sup>

$$\frac{F(a, b+1; c+1; z)}{F(a, b; c; z)} = \frac{1}{1 - \frac{a(c-b)z}{c(c+1)} \frac{1 - (b+1)(c-a+1)z}{(c+1)(c+2)} \frac{1 - (a+1)(c-b+1)z}{(c+2)(c+3)} \frac{1 - (b+2)(c-a+2)z}{(c+3)(c+4)} \dots} \quad (10)$$

The Gauss continued fraction converges everywhere in the complex plane, except on the real axis from +1 to  $+\infty$  and at isolated zeros of  $F(a, b; c; z)$ .

In order to obtain  $Q_v^m(\cos\theta + i0)$  from its continued fraction, we utilize the following Wronskian relations<sup>2</sup>:

$$W\{P_v^{-m}(z), Q_v^m(z)\} = P_v^{-m}(z) \frac{dQ_v^m(z)}{dz} - Q_v^m(z) \frac{dP_v^{-m}(z)}{dz} \quad (11)$$

and

$$W\{P_v^{-m}(z), Q_v^m(z)\} = \frac{e^{i\pi m}}{(1-z^2)}.$$

---

<sup>6</sup> H.S. Wall, *Analytic Theory of Continued Fractions*, d. Van Nostrand Company, Inc., New York, 1948.

Equating these, we have

$$P_v^{-m}(z) \frac{dQ_v^m(z)}{dz} - Q_v^m(z) \frac{dP_v^{-m}(z)}{dz} = \frac{e^{i\pi m}}{(1-z^2)} . \quad (12)$$

Now using the derivative relations<sup>1</sup>

$$(z^2-1) \frac{dP_v^{-m}(z)}{dz} = v z P_v^{-m}(z) - (v-m) P_{v-1}^{-m}(z) \quad (13)$$

and

$$(z^2-1) \frac{dQ_v^m(z)}{dz} = v z Q_v^m(z) - (v+m) Q_{v-1}^m(z) ,$$

we find

$$(v+m) P_v^{-m}(z) Q_{v-1}^m(z) - (v-m) P_{v-1}^{-m}(z) Q_v^m(z) = e^{i\pi m} . \quad (14)$$

Dividing both sides by  $Q_{v-1}^m(z)$  and replacing  $z$  by  $\cos\theta+i0$  , we get

$$(v+m) P_v^{-m}(\cos\theta+i0) - (v-m) P_{v-1}^{-m}(\cos\theta+i0) R_v^m(\cos\theta+i0) = \frac{e^{i\pi m}}{Q_{v-1}^m(\cos\theta+i0)} . \quad (15)$$

Replacing  $v$  by  $v+1$  and then solving for  $Q_v^m(\cos\theta+i0)$  ,

$$Q_v^m(\cos\theta+i0) = \frac{e^{i\pi m}}{[(v+m+1)P_{v+1}^{-m}(\cos\theta+i0) - (v-m+1)P_v^{-m}(\cos\theta+i0)R_{v+1}^m(\cos\theta+i0)]} . \quad (16)$$

Since<sup>1</sup>

$$P_{\nu}^{-m}(\cos\theta+i0) = e^{1/2 im\pi} P_{\nu}^{-m}(\cos\theta), \quad (17)$$

$$Q_{\nu}^m(\cos\theta+i0) = \frac{e^{1/2 im\pi}}{[(\nu+m+1)P_{\nu+1}^{-m}(\cos\theta) - (\nu-m+1)P_{\nu}^{-m}(\cos\theta)R_{\nu+1}^m(\cos\theta+i0)]}. \quad (18)$$

Now we can use the relation<sup>1</sup>

$$Q_{\nu}^m(\cos\theta) = e^{3/2 im\pi} Q_{\nu}^m(\cos\theta+i0) + i\pi/2 P_{\nu}^m(\cos\theta) \quad (19)$$

to finally get

$$Q_{\nu}^m(\cos\theta) = \frac{(-1)^m}{[(\nu+m+1)P_{\nu+1}^{-m}(\cos\theta) - (\nu-m+1)P_{\nu}^{-m}(\cos\theta)R_{\nu+1}^m(\cos\theta+i0)]} + i\pi/2 P_{\nu}^m(\cos\theta). \quad (20)$$

Results from programming the above method indicate that it works exceptionally well. Unfortunately, this method will likely run into difficulties for large  $|\nu|$ . Even though the Gauss continued fraction seems to converge more rapidly with increasing  $|\nu|$ , difficulties may be encountered because of losses in accuracy with  $P_{\nu}^m(\cos\theta)$ . For large  $|\nu|$ , it will be necessary to use asymptotic expansions to calculate  $Q_{\nu}^m(\cos\theta)$  and  $P_{\nu}^m(\cos\theta)$ .

Another region where trouble occurs is at small angles ( $\theta < 1^\circ$ ). The complex Gauss continued fraction algorithm is still very accurate at small angles, but the continued fraction converges at a very slow rate. Sometimes several thousand terms are required for convergence, which in turn uses an inordinate amount of computer time. This problem occurs because the argument approaches the cut of the Gauss continued fraction at  $+1$ . To avoid this problem, we use the logarithmic solution to calculate  $Q_{\nu}^m(\cos\theta)$  for small angles.

A similar method for computing  $Q_{\nu}^m(\cos\theta)$  has also been developed using real analysis only. Since no simple series formula for  $Q_{\nu}^m(\cos\theta)$  could be found, we concentrated on  $P_{\nu}^m(-\cos\theta)$  instead.  $P_{\nu}^m(-\cos\theta)$  is related to  $Q_{\nu}^m(\cos\theta)$  and  $P_{\nu}^m(\cos\theta)$  by<sup>1</sup>

$$Q_v^m(\cos\theta) = \frac{\pi}{2\sin[\pi(v+m)]} \{ \cos[\pi(v+m)] P_v^m(\cos\theta) - P_v^m(-\cos\theta) \}, \quad 0 < \theta < \pi/2. \quad (21)$$

Similar computational difficulties exist for the computation of  $P_v^m(-\cos\theta)$  as  $Q_v^m(\cos\theta)$ . Using the same equations as in calculating  $P_v^m(\cos\theta)$ , we can derive the following continued fraction

$$\frac{P_v^m(-\cos\theta)}{P_{v-1}^m(-\cos\theta)} = \frac{(v+m)}{(v-m)} \frac{F(-v, v+1; 1+m; (\cos\theta/2)^2)}{F(-v+1, v; 1+m; (\cos\theta/2)^2)}. \quad (22)$$

Letting  $a=-v$ ,  $b=v$ ,  $c=m$ ,  $z=(\cos\theta/2)^2$ , then

$$S_v^m(z) = \frac{F(a, b+1; c+1; z)}{F(a+1, b; c+1; z)} = \frac{F(a, b+1; c+1; z)}{F(a, b; c; z)} \frac{F(b, a; c; z)}{F(b, a+1; c+1; z)}. \quad (23)$$

This can be calculated by either two Gauss continued fractions or by two continued fractions of Frank<sup>7</sup> which are of the form

$$\frac{F(a, b; c; z)}{F(a, b+1; c+1; z)} =$$

(Equation is incomplete, continued on next page)

---

<sup>7</sup> E. Frank, "A New Class of Continued Fraction Expansions for the Ratios of Hypergeometric Functions," *Transactions of the American Mathematical Society*, 81, 454, 1956.

$$1 - \frac{a(c-b)z}{c(c+1)} + 1 - \frac{\frac{(a-b)z}{(c+1)}}{\frac{(a+1)(c+1-b)z}{(c+1)(c+2)}} + 1 - \frac{\frac{(a+1-b)z}{(c+2)}}{\frac{(a+2)(c+2-b)z}{(c+2)(c+3)}} + 1 - \dots \quad (24)$$

The continued fraction of Frank has its cut on the unit circle. It has proved to be of little value here because of its much slower convergence rate as compared to the Gauss continued fraction. This is especially true for small angles, which effectively limit the use of the Frank continued fraction to angles greater than  $30^\circ$ .

Utilizing similar Wronskian relations as in the complex analysis, we proceeded in a similar manner, with the exception of replacing  $Q_v^m(\cos\theta)$  by Eq. (21). The final result of the real analysis is then

$$Q_v^m(\cos\theta) = \frac{\pi}{2} P_v^m(\cos\theta) \cot[\pi(v+m)] + \frac{(-1)^m}{(v+1+m)[P_{v+1}^{-m}(\cos\theta) + P_v^{-m}(\cos\theta)S_{v+1}^m(z)]} \quad (25)$$

This result can easily be seen to be inferior to the complex analysis since it is undefined for integral  $v$ . Also, it is necessary to calculate two continued fractions which have relatively slow convergence rates which worsen with increasing  $|v|$ . Therefore, only the complex algorithm is used in the final programming. The real method will be useful in helping to check and compare the accuracy of the complex method for calculating  $Q_v^m(\cos\theta)$ .

## B. The Logarithmic Solution

As mentioned previously, the logarithmic solution is used for the calculation of  $Q_v^m(\cos\theta)$  at small angles ( $\theta < 1^\circ$ ). In this range, the

logarithmic solution is as accurate as the Gauss continued fraction method, but requires much less computer time to obtain results. From references [1] and [2], the following formulas can be derived.

$$\begin{aligned}
Q_v^m(\cos\theta) = & \frac{1}{2} P_v^m(\cos\theta) \left[ \log(1/\tan^2(\theta/2)) - 2\gamma - 2\psi(v+1) + \sum_{s=1}^m \frac{2s-1}{(v+s)(v+1-s)} \right] \\
& + \frac{(-1)^m}{2v} \left( \frac{1}{\tan\theta/2} \right)^m \sum_{r=0}^m (-v^2)(1-v^2)(4-v^2) \dots ((r-1)^2 - v^2)(v+r) \frac{(m-r-1)!}{r!} \\
& \times (-1)^r (\sin\theta/2)^{2r} \\
& + \frac{(\sin\theta/2)^m}{2v} \sum_{\ell=1}^{\infty} \frac{(-v^2)(1-v^2) \dots ((m+\ell-1)^2 - v^2)(m+v+\ell)}{\ell! (m+\ell)!} \\
& \times \left( 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{\ell} \right) (\sin\theta/2)^{2\ell} \\
& + \frac{(-1)^m}{2v} (v-m+1)(v-m+2) \dots (v+m) (\tan\theta/2)^m \\
& \times \sum_{r=0}^{\infty} \frac{(-v)^2(1-v^2) \dots ((r-1)^2 - v^2)(v+r)}{v! (r+m)!} \left( 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{m+r} \right) (\sin\theta/2)^{2r}
\end{aligned}$$

(26)

For  $m=0$ , but  $v$  not an integer,

$$Q_v^0(\cos\theta) = \frac{1}{2} P_v^0(\cos\theta) [\log(\cot^2(\theta/2)) - 2\gamma - 2\psi(v+1)]$$

$$+ \frac{1}{v} \sum_{\ell=1}^{\infty} \frac{(-v^2)(1-v^2)\dots((\ell-1)^2-v^2)(\ell+v)}{(\ell!)^2} \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{\ell}\right) (\sin\theta/2)^{2\ell} .$$

(26a)

For  $m=0$  and  $v=n$ , an integer,

$$Q_n^0(\cos\theta) = \frac{1}{2} P_n^0(\cos\theta) [\log(\cot^2(\theta/2)) - 2(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n})]$$

$$+ \sum_{\ell=1}^n \frac{(-1)^\ell (n+\ell)! (1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{\ell})}{(\ell!)^2 (n-\ell)!} (\sin\theta/2)^{2\ell} .$$

(26b)

#### IV. CONCLUSIONS

We have developed an accurate and efficient method of calculating the associated Legendre functions for integral order and complex degree. In general, the accuracy of  $P_v^m(\cos\theta)$  and  $Q_v^m(\cos\theta)$  was approximately equal to the maximum attainable with double precision arithmetic. By comparisons with special cases ( $P_{-1/2+i\lambda}^m(\cos\theta)$ ,  $Q_v^m(\cos\theta)$  for real  $v$ , Legendre Polynomials) and alternate means of calculation of the associated Legendre functions (Gaussian quadratures, alternate series), the modulus of the results appeared accurate



to at least 25 significant figures. Tabular and graphical results are presented in Appendix A and Appendix B, respectively.

In routines still under development at BRL, special series and nonhomogeneous recurrence formulas are being used to calculate  $\frac{\partial P_v^m(\cos\theta)}{\partial v}$  and  $\frac{\partial Q_v^m(\cos\theta)}{\partial v}$ . Also being programmed are asymptotic expansions to extend the range of  $|v|$  to very large values. For small angles ( $\theta < 30^\circ$ ), Bessel function expansions will be used.<sup>8</sup> Special hypergeometric series along with sines and cosines will be used for larger angles.

---

<sup>8</sup> N.K. Chu Khrukidze, "Asymptotic Formulae for Legendre Functions," USSR Computational Mathematics and Mathematical Physics, 6, 1966.

## REFERENCES

1. A. Erdelyi, et. al., Higher Transcendental Functions, Vol. 1, Bateman Manuscript Project, McGraw Hill Book Co., Inc., New York, 1953.
2. W. Magnus and F. Oberhettinger, Formulas and Theorems for the Special Functions of Mathematical Physics, Chelsea Publishing Co., New York, 1949.
3. M. Abramowitz and J.A. Stegun, Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, National Bureau of Standards, Washington, D.C., No. 55, Applied Mathematics Series, June 1964.
4. J.M. Smith, F.V.J. Olver and D.W. Lozier, "Legendre Polynomials," ACM Transactions on Mathematical Software, 7, 94-95, (1981).
5. A.S. Elder, "Formulas for Calculating Bessel Functions of Integral Order and Complex Argument," USA Ballistic Research Laboratory, Report No. BRL-TR-1423, 1968 AD# 680209.
6. H.S. Wall, Analytic Theory of Continued Fractions, d. Van Nostrand Company, Inc., New York, 1948.
7. E. Frank, "A New Class of Continued Fraction Expansions for the Ratios of Hypergeometric Functions," Transactions of the American Mathematical Society 81, 454, 1956.
8. N.K. Chu Khrukidze, "Asymptotic Formulae for Legendre Functions," USSR Computational Mathematics and Mathematical Physics, 6, 1966.

**APPENDIX A**  
**NUMERICAL CALCULATIONS**

LEGENDRE FUNCTION OF THE FIRST KIND  
COMPLEX DEGREE  
OF ORDER M=3

$P_3^M(x)$  (.5)

$T^c$	RE(P)	IM(P)	IP1
0.0	-.42625D+02	0.	.4262468784251533964540200D+02
5.0	-.49585D+02	.64943D+01	.5000834760959741973930324D+02
10.0	-.70211D+02	.52977D+01	.7041025119614440010311626D+02
15.0	-.10209D+03	-.13976D+02	.1030434935710465942931210D+03
20.0	-.13579D+03	-.64986D+02	.1505385830832665826182015D+03
25.0	-.14791D+03	-.15994D+03	.2178484845204532479989599D+03
30.0	-.96827D+02	-.29600D+03	.3114306679115888208824880D+03
35.0	.68674D+02	-.43319D+03	.4386022150293383826031352D+03
40.0	.37572D+03	-.47622D+03	.6065891000167963933163824D+03
45.0	.76870D+03	-.28849D+03	.8210502974015998485590649D+03
50.0	.10593D+04	.23040D+03	.1084084206793458916862205D+04
55.0	.95627D+03	.10115D+04	.1391983822476521931115219D+04
60.0	.23330D+03	.17175D+04	.1733287684730808667421165D+04
65.0	-.10143D+04	.18250D+04	.2087877903411993927468460D+04
70.0	-.22321D+04	.95508D+03	.2427864284513677923181747D+04
75.0	-.26230D+04	-.72251D+03	.2720658449418312029188117D+04
80.0	-.17002D+04	-.23912D+04	.2934000659324421154575291D+04
85.0	.22351D+03	-.30337D+04	.3041944970230033969349005D+04
90.0	.21091D+04	-.21758D+04	.3030254447274178917577798D+04

LEGENDRE FUNCTION OF THE FIRST KIND  
COMPLEX DEGREE  
OF ORDER M=10

$P_{10}^{10}(.5)$

$\tau^\circ$	RE (P)	IM (P)	!P!
0.0	-0.175670-23	0.	0.175674064520017855565315590-23
5.0	-0.457180+04	0.865330+04	0.97867218322653841041196340+04
10.0	-0.314130+05	0.176430+05	0.36028196115072803808138740+05
15.0	-0.112830+06	-0.170590+05	0.11411378337902771629119110+06
20.0	-0.208830+06	-0.253910+06	0.32876048346744076803377290+06
25.0	0.875990+05	-0.862090+06	0.86653132185731852923980780+06
30.0	0.169110+07	-0.124020+07	0.20970662619339281734298520+07
35.0	0.453820+07	0.112870+07	0.46764644832455623253170960+07
40.0	0.390000+07	0.881660+07	0.96406101840753922149059480+07
45.0	-0.921090+07	0.159540+08	0.18421790515796532619486960+08
50.0	-0.325880+08	0.265790+07	0.32695988141790905195968170+08
55.0	-0.332640+08	-0.425150+08	0.53981465789233646563728960+08
60.0	0.279720+08	-0.781350+08	0.82991034618931210835185680+08
65.0	0.116450+09	-0.239480+08	0.11888920593519715894629010+09
70.0	0.106550+09	0.117690+09	0.15876225186147215154304490+09
75.0	-0.608360+08	0.188070+09	0.19766507335310095814740260+09
80.0	-0.224820+09	0.459240+08	0.22946616538431742736989990+09
85.0	-0.162080+09	-0.188200+09	0.24837460556835351141106830+09
90.0	0.864590+08	-0.235270+09	0.25065147977222993928132580+09

LEGENDRE FUNCTION OF THE SECOND KIND  
COMPLEX DEGREE  
OF ORDER M=5

$Q_{5,0}^5(x)$  (.5)

T	RE (Q)	IM (Q)	!Q!
57.50	-.189660+05	-.279490+05	.33776251385318829792752810+05
57.55	-.547640+04	-.390930+05	.39474723994468911539028680+05
57.60	.144330+05	-.431730+05	.45521237126130919936229490+05
57.65	.367540+05	-.364910+05	.51792740815031983821134870+05
57.70	.553390+05	-.178050+05	.58133151683613064821378250+05
57.75	.635030+05	.104390+05	.6435538896355668994804760+05
57.80	.563310+05	.419700+05	.70247203980715377923883410+05
57.85	.329010+05	.680440+05	.75581268279396651032496650+05
57.90	-.255580+04	.800880+05	.80129267313383237769359590+05
57.95	-.412830+05	.727860+05	.83678895859810127956851070+05
58.00	-.724730+05	.463970+05	.86051872107269658884209020+05
58.05	-.868200+05	.722680+04	.87120589428358254946585150+05
58.10	-.798310+05	-.341310+05	.86820962370953959369312920+05
58.15	-.535050+05	-.662520+05	.85159449722613237042031150+05
58.20	-.156830+05	-.807030+05	.82213087359659378219558800+05
58.25	.227240+05	-.747450+05	.78122466852189489111992370+05
58.30	.514720+05	-.518760+05	.73078718534435249192152720+05
58.35	.642180+05	-.201540+05	.67306460096055501356699880+05
58.40	.600650+05	.108970+05	.61045167694305196344973680+05
58.45	.430490+05	.334740+05	.54531430237510680199921980+05
58.50	.200500+05	.435940+05	.47984093324010848496556470+05

LEGENDRE FUNCTION OF THE SECOND KIND  
COMPLEX DEGREE  
OF ORDER M=10

$$Q_5^0 e^{\tau} (.5)$$

T°	RE (Q)	IM (Q)	!Q!
57.50	-.30838D+08	.80986D+08	.8665892504808939120011191D+08
57.55	-.76502D+08	.42454D+08	.8749216251189213734382323D+08
57.60	-.97126D+08	-.30047D+08	.1016676631684915507256846D+09
57.65	-.67013D+08	-.11757D+09	.1353240616931585051295603D+09
57.70	.23620D+08	-.18191D+09	.1834338145301002362238196D+09
57.75	.15454D+09	-.17993D+09	.2371848993989002236938305D+09
57.80	.27473D+09	-.87713D+08	.2883900964858638118524417D+09
57.85	.32117D+09	.78045D+08	.3305155451242633059538781D+09
57.90	.25152D+09	.25686D+09	.3595011393852248646233413D+09
57.95	.73399D+08	.36727D+09	.3745369642717652245844263D+09
58.00	-.14957D+09	.34743D+09	.3782595851276043292972753D+09
58.05	-.32340D+09	.19140D+09	.3757938315855598531210807D+09
58.10	-.37001D+09	-.41902D+08	.3723724995989596125389625D+09
58.15	-.26828D+09	-.25533D+09	.3703607755943606701169766D+09
58.20	-.65276D+08	-.36197D+09	.3678087514662628100563560D+09
58.25	.14912D+09	-.32752D+09	.3598710208624954539994170D+09
58.30	.28871D+09	-.18294D+09	.3417854866099812991461015D+09
58.35	.31121D+09	-.23110D+07	.3112221502702430241729608D+09
58.40	.23110D+09	.13795D+09	.2691424684710954185029470D+09
58.45	.10224D+09	.19421D+09	.2194804360932069069491771D+09
58.50	-.14857D+08	.16760D+09	.1682617546537580692969713D+09

LEGENDRE FUNCTION OF THE SECOND KIND  
REAL DEGREE  
OF ORDER M=5

$Q_5^M$  (.5)

V	RE (Q)	IM (Q)	!Q!
.10	-.189010+03	-.213310-26	.18900801878022575115073880+03
.50	-.195440+03	-.512440-26	.19544238489553037926126100+03
1.00	-.209390+03	-.341670-26	.20938569762610427726198460+03
1.50	-.233880+03	-.132780-25	.23387516642480665348073920+03
2.00	-.270970+03	-.521710-26	.27096972633966435880962720+03
2.50	-.321050+03	.636140-26	.32105006758853366329468730+03
3.00	-.394140+03	-.978920-26	.39413778376678452190491220+03
5.00	-.185020+04	.297270-23	.18502430608177084179696150+04
10.00	.296550+04	-.335840-21	.29654539432231117608705680+04
15.00	.284100+06	-.502930-21	.28409853164708937102225370+06
20.00	.603750+06	.828500-20	.60375371292282375720737980+06
25.00	-.135870+07	.253050-19	.13587387943672703353943740+07
30.00	-.634910+07	-.238890-20	.63490511712187315497813710+07
40.00	.130790+08	-.368460-18	.13078752708656431332812410+08
50.00	.243290+08	.122650-17	.24328639125798572084814930+08



LEGENDRE FUNCTION OF THE SECOND KIND  
REAL DEGREE  
OF ORDER M=10

$Q_V^{(10)}$  (.5)

V	RE (Q)	IM (Q)	!Q!
.10	.442200+08	.786010-21	.44220194410630840630270270+08
.50	.449920+08	.839150-21	.44991529493579425711728760+08
1.00	.465400+08	-.113640-20	.465402311111111111111111+08
1.50	.488050+08	.588310-21	.48805089200901219389123100+08
2.00	.518960+08	.261400-20	.518963200000000000000000+08
2.50	.559720+08	.105640-20	.55972321436581460920884630+08
3.00	.612490+08	.151270-20	.612488533333333333333333+08
5.00	.102450+09	.361580-20	.102454613333333333333333+09
10.00	.193250+10	-.165340-17	.19325053906530602653438280+10
15.00	-.143380+12	.327860-15	.14337812638034767034623010+12
20.00	-.553540+12	-.236480-13	.55353879429590076544051080+12
25.00	.231670+14	-.879300-13	.23167488916661974139666970+14
30.00	.958260+14	.146730-11	.95826038452083520625193670+14
40.00	-.231000+16	-.265740-11	.23099684194763414252678960+16
50.00	.129730+17	-.281940-09	.12972832704690443884185730+17

CONICAL LEGENDRE FUNCTION OF THE FIRST KIND  
COMPLEX DEGREE  
OF ORDER M=1

$$P'_{-1/2} + iY(.5)$$

Y	RE (P)	IM (P)	!PI
.1	.155520+00	-.613440-37	.15552007267513654263023250+00
.5	.308900+00	-.788610-47	.30889931143353596441863850+00
1.0	.852490+00	-.717660-47	.85249333434783483823985720+00
1.5	.199730+01	.466350-35	.19973262751681190295872210+01
2.0	.418770+01	0.	.41877084758316896827828910+01
2.5	.824640+01	-.644430-32	.82464104821428748796918980+01
3.0	.156510+02	.380050-33	.15651379836797780973855520+02
5.0	.171430+03	-.732040-30	.17143494782052233848947420+03
10.0	.467810+05	-.823390-27	.46781022940599679876805790+05
15.0	.108530+08	-.126990-25	.10852960776347177009844210+08
20.0	.236410+10	-.263930-21	.23640638837358112648967540+10
25.0	.497810+12	-.208280-19	.49781366889965494786618090+12
30.0	.102630+15	.899050-17	.10262960807376627389172410+15
40.0	.419250+19	.818360-12	.41925103953819107803860260+19
50.0	.165700+24	-.354510-07	.16570401429028412573707120+24

CONICAL LEGENDRE FUNCTION OF THE FIRST KIND  
COMPLEX DEGREE  
OF ORDER M=10

$$P_{\frac{1}{2}+iy}^{(0)}(.5)$$

Y	RE(P)	IM(P)	!P!
.1	.489140+03	-.940100-26	.48913585134593560012170610+03
.5	.114780+04	-.293120-43	.11478410243829040410861450+04
1.0	.500700+04	-.421510-43	.50070386186075993238050600+04
1.5	.218580+05	.510360-31	.21857825018393532089200930+05
2.0	.920750+05	0.	.92075211208276866556112640+05
2.5	.373930+06	-.292220-27	.37392953232534501245459720+06
3.0	.146520+07	.355790-28	.14652154098920242714762300+07
5.0	.246960+09	-.105450-23	.24695549286117219972770460+09
10.0	.137230+14	-.241540-18	.13723468018111848115909750+14
15.0	.128830+18	-.150740-15	.12882834633792463793109230+18
20.0	.433690+21	.738980-08	.43369083034979352196541640+21
25.0	.779040+24	-.129130-04	.77904260498089933257409090+24
30.0	.925260+27	.223460-01	.92526360969238511960080180+27
40.0	.590890+33	-.885940+04	.59089366337138470757923660+33
50.0	.193830+39	.112640+10	.19382736426993325063842310+39

CONICAL LEGENDRE FUNCTION OF THE SECOND KIND  
COMPLEX DEGREE  
OF ORDER M=1

$$Q_2^{(1)} - iY^{(1)}(0.5)$$

Y	RE(Q)	IM(Q)	!Q!
.1	-.772640+00	-.743170-01	.77620144494795149481902260+00
.5	-.701770+00	-.445020+00	.83097629350916298288318260+00
1.0	-.540690+00	-.133410+01	.14395050700718725453807540+01
1.5	-.380260+00	-.313690+01	.31598501337947940203914340+01
2.0	-.255130+00	-.657800+01	.65829369379674706425735450+01
2.5	-.166710+00	-.129530+02	.12954500119931210979358670+02
3.0	-.107120+00	-.245850+02	.24585363008523722596962800+02
5.0	-.166620-01	-.269290+03	.26928938683620172809616360+03
10.0	-.123100-03	-.734830+05	.73483458998801770445147940+05
15.0	-.615430-07	-.170480+08	.17047790922334920422262420+08
20.0	.523280+01	-.371350+10	.37134628625998169909647760+10
25.0	.445490+08	-.781940+12	.78194189502694445116166370+12
30.0	.865310+14	.649100+14	.10817048658037239681947290+15
40.0	.498040+10	.658560+19	.65855799268166945812382480+19
50.0	.844200+05	.260290+24	.26028725698234736290819950+24

CONICAL LEGENDRE FUNCTION OF THE SECOND KIND  
COMPLEX DEGREE  
OF ORDER M=10

$$Q^{-1/2}_M(\gamma(.5))$$

Y	RE (Q)	IM (Q)	!Q!
.1	.437820+08	-.233740+03	.43781989399457307721149980+08
.5	.435000+08	-.165360+04	.43499831939445316981130020+08
1.0	.426310+08	-.783570+04	.42630825762273961890454350+08
1.5	.412240+08	-.343290+05	.41224355237761774941548300+08
2.0	.393400+08	-.144630+06	.39339978251706168803719860+08
2.5	.370540+08	-.587370+06	.37058171837997440309511730+08
3.0	.344540+08	-.230160+07	.34530799179481372592288350+08
5.0	.227690+08	-.387920+09	.38858442705555205139822480+09
10.0	.384040+07	-.215570+14	.21556773153737672893926180+14
15.0	.292870+06	-.202360+18	.20236309321467289828164950+18
20.0	-.700740+11	-.681240+21	.68123996330234816908619560+21
25.0	-.694020+19	-.122370+25	.12237197265179322867840160+25
30.0	-.996560+27	-.155210+28	.18444759122757150914669060+28
40.0	-.270780+25	.928170+33	.92817359724582902151311800+33
50.0	-.315580+21	.304460+39	.30446331182754753912411440+39

LEGENDRE FUNCTION OF THE SECOND KIND  
OF REAL DEGREE  $\nu=2$ .  
AND ORDER  $M=0$

$Q_2^c [\cos(T)]$

COMPLEX GAUSS  
CONTINUED FRACTION  
METHOD

SPECIAL CASE  
(FOR ACCURACY CHECK)  
 $Q_2^c [\cos(T)] = \text{LOG}[\cot^2(T/2)]$   
 $\times (3\cos^2(T)-1)/4 - 3/2\cos(T)$

$T^\circ$

.01	.7846543924828646523610051D+01	.7846543924828646523610051D+01
.10	.5543929083720490521528767D+01	.5543929083720490521528767D+01
.50	.3933951318124424740236797D+01	.3933951318124424740236797D+01
1.00	.3239410991461712058234597D+01	.3239410991461712058234597D+01
10.00	.8488417132323570536433152D+00	.8488417132323570536433152D+00
20.00	.213687158885539562955563D-01	.213687158885539562955563D-01
30.00	.4759394200986475272549308D+00	.4759394200986475272549308D+00
40.00	.7647683970726550081259554D+00	.7647683970726550081259554D+00
50.00	.8728124046175654413013493D+00	.8728124046175654413013493D+00
60.00	.8186632680417568557122028D+00	.8186632680417568557122028D+00
70.00	.6286869187008021321194748D+00	.6286869187008021321194748D+00
80.00	.3402505773017355382714222D+00	.3402505773017355382714222D+00

LEGENDRE FUNCTION OF THE SECOND KIND  
REAL DEGREE  
OF ORDER M=0

$Q_2^2(\cos(T))$

LOGARITHMIC  
SOLUTION

SPECIAL CASE  
(FOR ACCURACY CHECK)  
 $Q_2^2(\cos(T)) = \text{LOG}(\cos^2(T/2))$   
 $*(3\cos^2(T)-1)/4 - 3/2*\cos(T)$

T

.001	.1014912941946317922184000D+02	.1014912941946317922184000D+02
.005	.8539691402540814834167413D+01	.8539691402540814834167413D+01
.010	.7846543924828646523609732D+01	.7846543924828646523609732D+01
.050	.6237098088637495570857967D+01	.6237098088637495570857967D+01
.100	.5543929083720490521528448D+01	.5543929083720490521528448D+01
.250	.4627506207003646750005995D+01	.4627506207003646750005995D+01
.500	.3933951318124424740236479D+01	.3933951318124424740236479D+01
.750	.3527877952305230389869642D+01	.3527877952305230389869642D+01
1.000	.3239410991461712058234278D+01	.3239410991461712058234278D+01
5.000	.1601330656918337463046403D+01	.1601330656918337463046403D+01
10.000	.8488417132323570536429548D+00	.8488417132323570536429548D+00

LEGENDRE FUNCTION OF THE SECOND KIND  
OF DEGREE  $V=.1$   
OF ORDER  $M=1$

$Q_1^1(\cos(T))$

$T^\circ$	LOGARITHMIC SOLUTION	COMPLEX GAUSS CONTINUED FRACTION METHOD
.005	-.11459155967011203778918480+05	-.11459155967011203778918480+05
.010	-.57295780734439762339710660+04	-.57295780734439762339710660+04
.050	-.11459161236929236694631360+04	-.11459161236929236694631360+04
.100	-.57295879545620491117545330+03	-.57295879545620491117545330+03
.250	-.22918539897471327138320970+03	-.22918539897471327138320970+03
.500	-.11459578820359369037168810+03	-.11459578820359369037168810+03
.750	-.76400424582515658591572750+02	-.76400424582515658591572750+02
1.000	-.57303572639238578099893150+02	-.57303572639238578099893150+02
5.000	-.11490416444659423566983750+02	-.11490416444659423566983750+02
10.000	-.57855554431891268742160510+01	-.57855554431891268742160510+01



LEGENDRE FUNCTION OF THE SECOND KIND  
OF DEGREE  $\nu=10.5$   
OF ORDER  $M=1$

$Q_{10.5}^1(\cos(T))$

LOGARITHMIC  
SOLUTION

COMPLEX GAUSS  
CONTINUED FRACTION  
METHOD


$T^0$

.005	-.11459195770945768881171870+05	-.11459195770945768881171870+05
.010	-.57296503839534387814029990+04	-.57296503839534387814029990+04
.050	-.11461929533880912255798930+04	-.11461929533880912255798930+04
.100	-.57343946311317633105893220+03	-.57343946311317633105893220+03
.250	-.23014563670988900106772980+03	-.23014563670988900106772980+03
.500	-.11614982435180295584522110+03	-.11614982435180295584522110+03
.750	-.78407963278666147796055460+02	-.78407963278666147796055460+02
1.000	-.59671851253427766672521210+02	-.59671851253427766672521210+02
5.000	-.14122993109606857135954470+02	-.14122993109606857135954470+02
10.000	-.26651079459708560980737880+01	-.26651079459708560980737880+01


## **APPENDIX B**

### **GRAPHS**


$P_0(\cdot, 5)$  VS.  $V$




$Q_V(.5)$  VS.  $V$




$Q_1^{(1)}(x, y) \text{ vs. } x$




$P_1(.5)$  VS.  $V$



$P_3^3[\cos(\tau) + j \sin(\tau)](5) \text{ VS. } \tau^\circ$




$Q^2_{\{\cos(T)-isin(\eta)\}}(5)$  VS.  $T^\circ$






$P_{-1.5+i\gamma}^3(.5)$  VS.  $\gamma$



$Q^2_{s+i\gamma}(.5)$  VS  $\gamma$



# DISTRIBUTION LIST

<u>No. of</u> <u>Copies</u>	<u>Organization</u>	<u>No. of</u> <u>Copies</u>	<u>Organization</u>
12	Administrator Defense Technical Info Center ATTN: DTIC-DDA Cameron Station Alexandria, VA 22314	1	Commander USA AMCCOM ATTN: DRSMC-TDC(D) Dover, NJ 07801
1	Director Defense Advanced Research Projects Agency 1400 Wilson Boulevard Arlington, VA 22209	2	Commander USA AMCCOM ATTN: DRSMC-SC(D), B. Shulman Mr. Webster Dover, NJ 07801
1	Director Defense Nuclear Agency Washington, DC 20305	7	Commander USA AMCCOM ATTN: DRSMC-LCR-R(D), T. Moore DRSMC-LCR(D), W. Williver DRSMC-LCS-D(D), K. Rubin DRSMC-LCA(D), T. Davidson DRSMC-LCU-CT(D), E. Barrieres DRSMC-LCA-G(D), J. Lannon DRSMC-LCU-CV(D), E. Fennel Dover, NJ 07801
1	Headquarters Department of the Army ATTN: DAMA-WSW, LTC Ira Dorsey Washington, DC 20310	2	Commander USA AMCCOM ATTN: DRSMC-SE(D) Dover, NJ 07801
1	Director USA BMD Advanced Technology Center ATTN: BMDATC-M P.O. Box 1500 Huntsville, AL 35807	1	Commander USA AMCCOM ATTN: DRSMC-FU(D) Dover, NJ 07801
1	Commander USA Materiel Development and Readiness Command ATTN: DRCMD-ST 5001 Eisenhower Avenue Alexandria, VA 22333	2	Commander USA AMCCOM ATTN: DRSMC-LC(D) DRSMC-TDR(D) Dover, NJ 07801
1	Commander USA Materiel Development and Readiness Command ATTN: DRCDE-SG, G. Duncan 5001 Eisenhower Avenue Alexandria, VA 22333	2	Commander USA AMCCOM ATTN: DRSMC-QAT-B(D), W. Denery Dover, NJ 07801
2	Commander USA AMCCOM ATTN: DRSMC-TSS(D) Dover, NJ 07801		

# DISTRIBUTION LIST

<u>No. of</u> <u>Copies</u>	<u>Organization</u>	<u>No. of</u> <u>Copies</u>	<u>Organization</u>
1	Commander USA AMCCOM ATTN: DRSMC-PRW-C(D), F. Baldinger Dover, NJ 07801	1	Director USA Air Mobility Research and Development Laboratory Ames Research Center Moffett Field, CA 94035
1	Commander USA AMCCOM ATTN: DRSMC-LMC(D), MAJ Wolf Dover, NJ 07801	1	Commander USA Communications Research and Development Command ATTN: DRSEL-ATDD Fort Monmouth, NJ 07703
2	Commandant USA Infantry School ATTN: ATSH-CD-CSO-OR Fort Benning, GA 31905	1	Commander USA Electronics Research and Development Command Technical Support Activity ATTN: DELSD-L Fort Monmouth, NJ 07703
1	Commander USA Armament, Munitions & Chem Cmd ATTN: DRSMC-LEP-L(R) Rock Island, IL 61299	1	Commander USA Harry Diamond Labs 2800 Powder Mill Road Adelphi, MD 20783
2	Director USA AMCCOM Benet Weapons Laboratory ATTN: DRSMC-LCB-TL(D) DRSMC-LCB-DR(D), W. Jarrett Watervliet, NY 12189	1	Commander USA Missile Command ATTN: DRSMI-R Redstone Arsenal, AL 35898
2	Director USA AMCCOM Benet Weapons Laboratory ATTN: DRSMC-LCB(D), T. Simkins Watervliet, NY 12189	1	Commander USA Missile Command ATTN: DRSMI-YDL Redstone Arsenal, AL 35898
1	Commander USA Aviation Research and Development Command ATTN: DRDAV-E 4300 Goodfellow Blvd St. Louis, MO 63120	1	Commander USA Mobility Equipment Research & Development Command Fort Belvoir, VA 22060
2	Director US Army Research and Technology Laboratories (AVRADCOM) Ames Research Center Moffett Field, CA 94035	2	Commander USA Tank Automotive Command ATTN: DRSTA-TSL Warren, MI 48090
		1	Commander USA Tank Automotive Command ATTN: DRCPM-PG, COL J. Welsh Warren, MI 48090

# DISTRIBUTION LIST

<u>No. of</u> <u>Copies</u>	<u>Organization</u>	<u>No. of</u> <u>Copies</u>	<u>Organization</u>
1	Commander USA Tank Automotive Command ATTN: DRCPM-LA, COL. B. McClain Warren, MI 48090	1	Director USA TRADOC Systems Analysis Activity ATTN: ATAA-SL White Sands Missile Range NM 88002
1	Director TRASANA ATTN: ATOR-TFB, R. Reynolds White Sands Missile Range NM 88002	1	Commandant USA Infantry School ATTN: ATSH-CD Ft. Benning, GA 31905
2	Commander USA Armor Center ATTN: ATZK-CD-MS, COL G. Duncan ATZK-CD-MS, LTC R. Ellis Ft. Knox, KY 40121	1	Commander Fort Monroe ATTN: ATCD-ML Hampton, VA 23365
1	Director of Combat Development USA Armor Center ATTN: ATZK-CD, COL Demont Ft. Knox, KY 40121	1	Commander 9th Infantry Division ATTN: AFVO-TB-MS, COL J. Ott Ft. Lewis, WA 98433
3	Commander USA Research Office P.O. Box 12211 ATTN: Technical Director Engineering Division Metallurgy & Materials Division Research Triangle Park NC 27709	1	Commander USA Combined Arms Combat Development Activity ATTN: ATZL-CAD-IC, MAJ M. Cocshignano Ft. Leavenworth, KS 66027
2	Commander USA Research Office ATTN: J. Chandra F. Schmiedeshoff Research Triangle Park NC 27709	3	Commander Naval Sea Systems Command ATTN: SEA-62R, SEA-62Y, SEA-9961 Washington, DC 20360
4	Director USA Materials and Mechanics Research Center ATTN: Director (3 cys) DRXMR-ATL (1 cy) Watertown, MA 02172	1	Commander David W. Taylor Naval Ship Research and Development Center Bethesda, MD 20084
		1	Commander Naval Air Systems Command Washington, DC 20360

# DISTRIBUTION LIST

<u>No. of Copies</u>	<u>Organization</u>	<u>No. of Copies</u>	<u>Organization</u>
1	Commander Naval Air Development Center, Johnsville Warminster, PA 18974	1	Commander US Marine Corps Development and Education Command ATTN: Army Liaison, COL C. Beaumont Quantico, VA 22134
1	Commander Naval Missile Center Point Mugu, CA 93041	1	Commander US Marine Corps Development and Education Command ATTN: D093, LTC T. Kelly Quantico, VA 22134
1	Commander Naval Surface Weapons Center ATTN: G31, E. Wilkerson Dahlgren, VA 22448	1	Project Manager Division Air Defense Gun ATTN: DRCPM-ADG Dover, NJ 07801
1	Commander Naval Surface Weapons Center Silver Spring, MD 20910	3	Project Manager Cannon Artillery Weapons System ATTN: DRCPM-CAWS Dover, NJ 07801
1	Commander Naval Weapons Center China Lake, CA 93555	2	Project Manager Nuclear Munitions ATTN: DRCPM-NUC Dover, NJ 07801
1	Commander Naval Research Laboratory Washington, DC 20375	2	Project Manager Tank Main Armament Systems ATTN: DRCPM-TMA Dover, NJ 07801
1	Superintendent Naval Postgraduate School ATTN: Dir of Lib Monterey, CA 93940	1	Product Manager for 30mm Ammo ATTN: DRCPM-AAH-30mm Dover, NJ 07801
1	Commander Naval Ordnance Station Indian Head, MD 20640	2	Product Manager M110E2 Weapon System, DARCOM ATTN: DRCPM-M110E2 Rock Island, IL 61299
1	Commandant US Marine Corps ATTN: RDS, LTC H. May Washington, DC 20380		
1	Commander US Marine Corps Development and Education Command ATTN: D093, COL J. Molineaux Quantico, VA 22134		

# DISTRIBUTION LIST

<u>No. of</u> <u>Copies</u>	<u>Organization</u>	<u>No. of</u> <u>Copies</u>	<u>Organization</u>
1	Director National Aeronautics and Space Administration Langley Research Center Hampton, VA 23365	2	Commander US Army Jefferson Proving Ground ATTN: STEJP-TD-O, A. Tilley STEJP-TD-E, J. Tooney Madison, IN 47251
1	Director National Aeronautics and Space Administration Manned Spacecraft Center ATTN: Lib Houston, TX 77058	2	Commander US Army Yuma Proving Ground ATTN: STEYP-MTW, R. Torp G. Stullenbarger Yuma, AZ 85364
2	AFRPL ATTN: W. Andrepont T. Park Edwards AFB, CA 93523	1	President US Army Armor and Engineer Board ATTN: ATZK-RE-IN, L. Smith Fort Knox, KY 40121
1	AFATL Eglin AFB, FL 32542	<u>Aberdeen Proving Ground</u>	
1	AFWL/SUL Kirkland AFB, NM 87117	Dir, USAMSAA ATTN: DRXSY-D DRXSY-MP, H. Cohen DRXSY-GA, B. Siegel DRXSY-GA, R. Conroy	
1	Director Los Alamos Scientific Lab P.O. Box 1663 Los Alamos, NM 87544	Dir, MTD ATTN: H. King P. Paules	
1	Director Lawrence Livermore Laboratory P.O. Box 808 Livermore, CA 94550	Cdr, USATECOM ATTN: DRSTE-TO-F DRSTE-CM-R, L. Saubier	
1	BLM Applied Mechanics Consultants ATTN: Dr. A. Boresi 3310 Willett Drive Laramie, WY 82070	Cdr, CRDC, AMCCOM ATTN: DRSMC-CLB-PA DRSMC-CLN DRSMC-CLJ-L	
1	AAI Corporation ATTN: J. Hebert P.O. Box 126 Cockeysville, MD 21030		
1	ARES, Inc. ATTN: W. Ford 818 Front Street Port Clinton, OH 43452		

## USER EVALUATION OF REPORT

Please take a few minutes to answer the questions below; tear out this sheet, fold as indicated, staple or tape closed, and place in the mail. Your comments will provide us with information for improving future reports.

1. BRL Report Number \_\_\_\_\_

2. Does this report satisfy a need? (Comment on purpose, related project, or other area of interest for which report will be used.)

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

3. How, specifically, is the report being used? (Information source, design data or procedure, management procedure, source of ideas, etc.) \_\_\_\_\_

\_\_\_\_\_  
\_\_\_\_\_

4. Has the information in this report led to any quantitative savings as far as man-hours/contract dollars saved, operating costs avoided, efficiencies achieved, etc.? If so, please elaborate.

\_\_\_\_\_  
\_\_\_\_\_

5. General Comments (Indicate what you think should be changed to make this report and future reports of this type more responsive to your needs, more usable, improve readability, etc.) \_\_\_\_\_

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

6. If you would like to be contacted by the personnel who prepared this report to raise specific questions or discuss the topic, please fill in the following information.

Name: \_\_\_\_\_

Telephone Number: \_\_\_\_\_

Organization Address: \_\_\_\_\_

\_\_\_\_\_  
\_\_\_\_\_